

On conserved quantities in the theory of charged boson fields of spin 0 and 1

Tokarevskaya N.G., Red'kov V.M.¹

Institute of Physics, National Academy of Sciences of Belarus
68 Skarina str., Minsk, BELARUS

Bosons of spin 0 and 1, with different intrinsic parities, are described by full sets of spinor equations in the frame of the Dirac-Kahler theory. This enables us to obtain the conservation laws for the boson particles with one value of spin by imposing linear additional conditions in the known sixteen conserved currents of the Dirac-Kahler field. In this way for each boson the known conserved quantities, charge vector $j^a(x)$, symmetrical energy-momentum tensor $T^{ab}(x)$, and angular moment tensor $L^{a[bc]}(x)$, have been found. Additionally, for scalar fields, one conserved current $\nu^a(x)$ has been constructed; it is not zero one only for a complex-valued field. For a vector particles, two additional currents, $\nu^a(x)$ and $\nu^{ab}(x)$, are found that again do not vanish when fields are complex-valued. Those currents $\nu^a(x)$ and $\nu^{ab}(x)$ have not seemingly any physical interpretation.

1 Introduction

It is known that the wave equation of the Dirac-Kahler field allows for sixteen conservation laws and there exists a number of works where such conserved Dirac-Kahler currents have been examined (for example, see in [1-19]). However previously, as a rule, the main attention was given to existence of those conservation laws and description of its underlying symmetry instead of elucidating its physical meaning. Else one point significant for us is that the Dirac-Kahler theory gives possibility to obtain many facts of the more simple theory of ordinary boson particles of spin 0 or 1 ; this is achieved by imposing special linear conditions on 16-component Dirac-Kahler wave function [20]. The same trick may be done with the Dirac-Kahler conserved currents, and it is the object of the present note.

2 Conserved quantities in the theory of the Dirac-Kahler field

Before proceeding to the problem of conserved quantities in the Dirac-Kahler theory, with subsequent analysis of the most interest cases of ordinary bosons particles of spin 0 or 1 and different intrinsic parity, let us consider in the beginning the simple example of the Dirac particle – at this there exists possibility to introduce the notation required.

The Dirac equation and its conjugate for $\bar{\Psi}(x) = \Psi^+(x)\gamma^0$ have the form

$$[i\gamma^a \vec{\partial}_a - m] \Psi(x) = 0 , \quad \bar{\Psi}(x) [i\gamma^a \overleftarrow{\partial}_a + m] = 0 . \quad (1)$$

¹E-mail: redkov@dragon.bas-net.by

In the following we are to take the formulas (the use of spinor basis is presupposed)

$$(\gamma^0)^+ = +\gamma^0, \quad (\gamma^i)^+ = -\gamma^i, \quad \gamma^0(\gamma^a)^+\gamma^0 = +\gamma^a. \quad (2)$$

From (1) through the known procedure it follows the charge conservation law

$$\partial_a J^a(x) = 0, \quad J^a(x) = \bar{\Psi}(x)\gamma^a\Psi(x). \quad (3)$$

The symmetry properties of the vector $J^a(x)$ under double spin covering of the full Lorentz group G_L^{spin}

$$S(k, \bar{k}^*), \quad M = i\gamma^0, \quad N = \gamma^0\gamma^5$$

can be easily described:

$$\begin{aligned} S(k, \bar{k}^*) : \quad & J'^a(x') = L^a{}_b(k, k^*) J^b(x), \\ M : \quad & J'^a(x') = +L^{(P)a}{}_b J^b(x), \\ N : \quad & J'^a(x') = -L^{(T)a}{}_b J^b(x). \end{aligned} \quad (4)$$

The problem of conserved currents in the Dirac-Kahler theory may be considered in much the same manner. The main equation and its conjugate are

$$[i\gamma^a \vec{\partial}_a - m] U(x) = 0, \quad \bar{U}(x) [i\gamma^a \overleftarrow{\partial}_a + m] = 0, \quad (5)$$

$$\bar{U}(x) = U^+(x) (\gamma^0 \otimes \gamma^0) = \tilde{\gamma}^0 U^+(x) \gamma^0. \quad (6)$$

The Dirac-Kahler functions behave under the group G_L^{spin} as follows:

$$\begin{aligned} S(k, \bar{k}^*) : \quad & U'(x') = (S \otimes S) U(x) = S U(x) \tilde{S}, \\ & \bar{U}'(x') = \bar{U}(x) (S^{-1} \otimes S^{-1}) = \tilde{S}^{-1} \bar{U}(x) S^{-1}, \end{aligned} \quad (7)$$

$$\begin{aligned} M = i\gamma^0 : \quad & U'(x') = (M \otimes M) U(x) = M U(x) \tilde{M}, \\ & \bar{U}'(x') = \bar{U}(x) (M^{-1} \otimes M^{-1}) = \tilde{M}^{-1} \bar{U}(x) M^{-1}, \end{aligned} \quad (8)$$

and

$$\begin{aligned} N = \gamma^0\gamma^5 : \quad & U'(x') = (N \otimes N) U(x) = N U(x) \tilde{N}, \\ & \bar{U}'(x') = \bar{U}(x) (N^{-1} \otimes N^{-1}) = \tilde{N}^{-1} \bar{U}(x) N^{-1}. \end{aligned} \quad (9)$$

Combining equations from (5), we get to sixteen conserved currents:

$$\begin{aligned} \bar{U} [i\gamma^a \vec{\partial}_a - m] U(x) + \bar{U}(x) [i\gamma^a \overleftarrow{\partial}_a + m] U(x) = 0 : \quad & \implies \\ \partial_a J^a(x) = 0, \quad \text{where} \quad & J^a(x) = \bar{U}(x) \gamma^a U(x). \end{aligned} \quad (10)$$

Take notice that $J^a(x)$ is a 4×4 matrix; that is it represents 16 conserved quantities. On the base of these one may construct special linear combinations which will be of simple symmetry properties under the group G_L^{spin} :

$$\begin{aligned}
j^{(a)}(x) &= \text{Sp } J^a(x) , & \partial_a j^{(a)}(x) &= 0 ; \\
\nu^{(a)}(x) &= \text{Sp } \gamma^5 J^a(x) , & \partial_a \nu^{(a)}(x) &= 0 ; \\
T^{(a)l}(x) &= \text{Sp } \tilde{\gamma}^l J^a(x) , & \partial_a T^{(a)l}(x) &= 0 ; \\
\nu^{(a)l}(x) &= \text{Sp } \gamma^5 \tilde{\gamma}^l J^a(x) , & \partial_a \nu^{(a)l}(x) &= 0 ; \\
L^{(a)kl}(x) &= \text{Sp } \tilde{\sigma}^{kl} J^a(x) , & \partial_a \varphi^{(a)kl}(x) &= 0 .
\end{aligned} \tag{11}$$

Now the task is to follow the symmetry properties of these separate currents under the group G_L^{spin} . First, let us consider the current $j^{(a)}(x)$:

$$\begin{aligned}
S(k, \bar{k}^*) : & \quad j^{('a)} = L^a{}_b(k, , k^*) j^{(b)} , \\
M : & \quad j^{('a)} = L^{(P)a}{}_b j^{(b)} , \\
N : & \quad j^{('a)} = L^{(T)a}{}_b j^{(b)} .
\end{aligned} \tag{12}$$

For the current $\nu^{(a)}(x)$

$$\begin{aligned}
S(k, \bar{k}^*) : & \quad \nu^{('a)}(x') = +L^a{}_b(k, , k^*) \nu^{(b)}(x) , \\
M : & \quad \nu^{('a)}(x') = -L^{(P)a}{}_b \nu^{(b)}(x) , \\
N : & \quad \nu^{('a)}(x') = -L^{(T)a}{}_b \nu^{(b)}(x) .
\end{aligned} \tag{13}$$

For the current $T^{(a)l}(x)$

$$\begin{aligned}
S(k, \bar{k}^*) : & \quad T^{('a)l}(x') = L^l{}_k L^a{}_b T^{(b)k}(x) , \\
M : & \quad T^{('a)l}(x') = L^{(P)l}{}_k L^{(P)a}{}_b T^{(b)k}(x) , \\
N : & \quad T^{('a)l}(x') = L^{(T)l}{}_k L^{(T)a}{}_b T^{(b)k}(x) .
\end{aligned} \tag{14}$$

For the current $\nu^{(a)l}(x)$

$$\begin{aligned}
S(k, \bar{k}^*) : & \quad \nu^{('a)l}(x') = L^l{}_k L^a{}_b \nu^{(b)k}(x) , \\
M : & \quad \nu^{('a)l}(x') = -L^{(P)l}{}_k L^{(P)a}{}_b \nu^{(b)k}(x) , \\
N : & \quad \nu^{('a)l}(x') = -L^{(T)l}{}_k L^{(T)a}{}_b \nu^{(b)k}(x) .
\end{aligned} \tag{15}$$

And for the current $L'^{(a)bc}(x')$ we have

$$\begin{aligned}
S(k, \bar{k}^*) : & \quad L'^{(a)bc}(x') = L^a{}_b L^b{}_k L^c{}_l L^{(b)kl}(x) , \\
M : & \quad L'^{(a)bc}(x') = L^{(P)a}{}_b L^{(P)b}{}_k L^{(P)c}{}_l L^{(b)kl}(x) , \\
N : & \quad L'^{(a)bc}(x') = L^{(T)a}{}_b L^{(T)b}{}_k L^{(T)c}{}_l L^{(b)kl}(x) .
\end{aligned} \tag{16}$$

3 Conserved currents for boson fields in tensor form

Now the task is to obtain explicit tensor form for the above currents (11). With the use of the formula for the bispinor metric matrix

$$E = \begin{vmatrix} i\sigma^2 & 0 \\ 0 & -i\sigma^2 \end{vmatrix} = i \gamma^0 \gamma^2, \quad E^{-1} = -i \gamma^0 \gamma^2,$$

the Dirac-Kahler function looks as

$$U(x) = \left[-i A + \gamma^5 B + \gamma^l A_l + i \gamma^l \gamma^5 B_l + i \sigma^{mn} F_{mn} \right] (-i \gamma^0 \gamma^2), \quad (17)$$

also

$$U^+(x) = (i \gamma^0 \gamma^2) \left[+i A^* + \gamma^5 B^* + (\gamma^l)^+ A_l^* - i \gamma^5 (\gamma^l)^+ B_l^* - i (\sigma^{mn})^+ F_{mn}^* \right],$$

and further

$$\bar{U}(x) = (-i \gamma^0 \gamma^2) \left[+i A^* - \gamma^5 B^* + \gamma^l A_l^* + i \gamma^5 \gamma^l B_l^* + i \sigma^{mn} F_{mn}^* \right]. \quad (18)$$

Therefore, the conserved current matrix may be given as

$$\begin{aligned} J^{(a)}(x) &= \bar{U}(x) \gamma^a U(x) = \\ &= (-i \gamma^0 \gamma^2) \left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \\ &\times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) (-i \gamma^0 \gamma^2). \end{aligned} \quad (19)$$

Also we need two relations:

$$\tilde{\gamma}^l (-i \gamma^0 \gamma^2) = -\gamma^l (-i \gamma^0 \gamma^2), \quad \tilde{\sigma}^{ab} (-i \gamma^0 \gamma^2) = -(-i \gamma^0 \gamma^2) \sigma^{ab}.$$

So that, from (11) it follows the formulas for conserved currents of the Dirac-Kahler field:

$$\begin{aligned} j^{(a)}(x) &= \text{Sp } J^a(x) = \\ &= -\text{Sp} \left[\left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \right. \\ &\quad \left. \times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) \right], \end{aligned} \quad (20)$$

$$\begin{aligned} \nu^{(a)}(x) &= \text{Sp } \gamma^5 J^a(x) = \\ &= -\text{Sp} \left[\gamma^5 \left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \right. \\ &\quad \left. \times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) \right], \end{aligned} \quad (21)$$

$$\begin{aligned} T^{(a)l}(x) &= \text{Sp } \tilde{\gamma}^l J^a(x) = \\ &= +\text{Sp} \left[\gamma^l \left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \right. \\ &\quad \left. \times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) \right], \end{aligned} \quad (22)$$

$$\begin{aligned}
\nu^{(a)l}(x) &= \text{Sp } \gamma^5 \tilde{\gamma}^l J^a(x) = \\
&= + \text{Sp } \left[\gamma^5 \gamma^l \left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \right. \\
&\quad \left. \times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) \right] , \tag{23}
\end{aligned}$$

$$\begin{aligned}
L^{(a)kl}(x) &= \text{Sp } \tilde{\sigma}^{kl} J^a(x) = \\
&= + \text{Sp } \left[\sigma^{kl} \left(+i A^* - \gamma^5 B^* + \gamma^p A_p^* + i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^* \right) \times \right. \\
&\quad \left. \times \gamma^a \left(-i A + \gamma^5 B + \gamma^s A_s + i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn} \right) \right] . \tag{24}
\end{aligned}$$

The relationships presuppose rather unwieldy calculation. For us in the first place the most interesting are ordinary particles with one value of spin, 0 or 1, and certain intrinsic parity. So, instead of (3,4) we will get the four expansions for more simple cases $S = 0, \tilde{0}, 1, \tilde{1}$.

Particle $S = 0$

$$\begin{aligned}
j^{(a)}(x) &= - \text{Sp } \left[\left(+i A^* + \gamma^p A_p^* \right) \gamma^a \left(-i A + \gamma^s A_s \right) \right] , \\
\nu^{(a)}(x) &= - \text{Sp } \left[\gamma^5 \left(+i A^* + \gamma^p A_p^* \right) \gamma^a \left(-i A + \gamma^s A_s \right) \right] , \\
T^{(a)l}(x) &= + \text{Sp } \left[\gamma^l \left(+i A^* + \gamma^p A_p^* \right) \gamma^a \left(-i A + \gamma^s A_s \right) \right] , \tag{25} \\
\nu^{(a)l}(x) &= + \text{Sp } \left[\gamma^5 \gamma^l \left(+i A^* + \gamma^p A_p^* \right) \gamma^a \left(-i A + \gamma^s A_s \right) \right] , \\
L^{(a)kl}(x) &= + \text{Sp } \left[\sigma^{kl} \left(+i A^* + \gamma^p A_p^* \right) \gamma^a \left(-i A + \gamma^s A_s \right) \right] .
\end{aligned}$$

Particle $S = \tilde{0}$

$$\begin{aligned}
j^{(a)}(x) &= - \text{Sp } \left[\left(-\gamma^5 B^* + i \gamma^5 \gamma^p B_p^* \right) \gamma^a \left(+\gamma^5 B + i \gamma^s \gamma^5 B_s \right) \right] , \\
\nu^{(a)}(x) &= - \text{Sp } \left[\gamma^5 \left(-\gamma^5 B^* + i \gamma^5 \gamma^p B_p^* \right) \gamma^a \left(+\gamma^5 B + i \gamma^s \gamma^5 B_s \right) \right] , \\
T^{(a)l}(x) &= + \text{Sp } \left[\gamma^l \left(-\gamma^5 B^* + i \gamma^5 \gamma^p B_p^* \right) \gamma^a \left(+\gamma^5 B + i \gamma^s \gamma^5 B_s \right) \right] , \tag{26} \\
\nu^{(a)l}(x) &= + \text{Sp } \left[\gamma^5 \gamma^l \left(-\gamma^5 B^* + i \gamma^5 \gamma^p B_p^* \right) \gamma^a \left(+\gamma^5 B + i \gamma^s \gamma^5 B_s \right) \right] , \\
L^{(a)kl}(x) &= + \text{Sp } \left[\sigma^{kl} \left(-\gamma^5 B^* + i \gamma^5 \gamma^p B_p^* \right) \gamma^a \left(+\gamma^5 B + i \gamma^s \gamma^5 B_s \right) \right] .
\end{aligned}$$

Particle $S = 1$

$$\begin{aligned}
j^{(a)}(x) &= - \text{Sp } \left[\left(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^* \right) \gamma^a \left(+\gamma^s A_s + i \sigma^{mn} F_{mn} \right) \right] , \\
\nu^{(a)}(x) &= - \text{Sp } \left[\gamma^5 \left(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^* \right) \gamma^a \left(+\gamma^s A_s + i \sigma^{mn} F_{mn} \right) \right] , \\
T^{(a)l}(x) &= + \text{Sp } \left[\gamma^l \left(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^* \right) \gamma^a \left(+\gamma^s A_s + i \sigma^{mn} F_{mn} \right) \right] , \tag{27} \\
\nu^{(a)l}(x) &= + \text{Sp } \left[\gamma^5 \gamma^l \left(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^* \right) \gamma^a \left(+\gamma^s A_s + i \sigma^{mn} F_{mn} \right) \right] , \\
L^{(a)kl}(x) &= + \text{Sp } \left[\sigma^{kl} \left(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^* \right) \gamma^a \left(+\gamma^s A_s + i \sigma^{mn} F_{mn} \right) \right] .
\end{aligned}$$

Particle $S = \tilde{1}$

$$\begin{aligned}
j^{(a)}(x) &= - \text{Sp} \left[(+i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn}) \right] , \\
\nu^{(a)}(x) &= - \text{Sp} \left[\gamma^5 (+i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn}) \right] , \\
T^{(a)l}(x) &= + \text{Sp} \left[\gamma^l (+i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn}) \right] , \quad (28) \\
\nu^{(a)l}(x) &= + \text{Sp} \left[\gamma^5 \gamma^l (+i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn}) \right] , \\
L^{(a)kl}(x) &= + \text{Sp} \left[\sigma^{kl} (+i \gamma^5 \gamma^p B_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+i \gamma^s \gamma^5 B_s + i \sigma^{mn} F_{mn}) \right] .
\end{aligned}$$

First, let us consider the case of scalar particle $S = 0$, starting from the current $j^{(a)}(x)$ (take notice on factor 1/4)

$$S = 0 : \quad j^{(a)}(x) = - \frac{1}{4} \text{Sp} \left[(+i A^* + \gamma^p A_p^*) \gamma^a (-i A + \gamma^s A_s) \right] , \quad (29)$$

an auxiliary table (the all non-zero traces of matrix combinations)

$$\begin{array}{ccc}
& \gamma^a & \gamma^a \gamma^s \\
I & 0 & \text{Sp} (\gamma^a \gamma^s) \\
\gamma^p & \text{Sp} (\gamma^p \gamma^a) & 0
\end{array} ,$$

from this it follows

$$S = 0 : \quad j^{(a)}(x) = \frac{1}{i} (A^* A^a - A A^{a*}) . \quad (3.9c)$$

The current vanishes identically for a real-valued scalar field. Now in the same manner consider the current

$$S = 0 : \quad \nu^{(a)}(x) = - \frac{1}{4} \text{Sp} \left[\gamma^5 (+i A^* + \gamma^p A_p^*) \gamma^a (-i A + \gamma^s A_s +) \right] ; \quad (30)$$

an auxiliary table

$$\begin{array}{ccc}
& \gamma^a & \gamma^a \gamma^s \\
\gamma^5 & 0 & 0 \\
\gamma^5 \gamma^p & 0 & 0
\end{array} ,$$

so that

$$S = 0 : \quad \nu^{(a)}(x) \equiv 0 . \quad (31)$$

Consider the current

$$S = 0 : \quad T^{(a)l}(x) = + \frac{1}{4} \text{Sp} \left[\gamma^l (+i A^* + \gamma^p A_p^*) \gamma^a (-i A + \gamma^s A_s) \right] ; \quad (32)$$

an auxiliary table

$$\begin{array}{ccc} \gamma^l & \gamma^a & \gamma^a \gamma^s \\ \gamma^l \gamma^p & \text{Sp} (\gamma^l \gamma^a) & 0 \\ & 0 & \text{Sp} (\gamma^l \gamma^p \gamma^a \gamma^s) \end{array} ,$$

non-zero traces

$$\frac{1}{4} \text{Sp} (\gamma^l \gamma^a) = g^{la} , \quad \frac{1}{4} \text{Sp} (\gamma^l \gamma^p \gamma^a \gamma^s) = g^{lp} g^{as} - g^{la} g^{ps} + g^{ls} g^{pa} ,$$

the conserved current is

$$S = 0 : \quad T^{(a)l}(x) = A^* A g^{al} - g^{al} (A_s^* A^s) + (A^{a*} A^l + A^a A^{l*}) . \quad (33)$$

This quantity differs from zero both for complex- and real-valued scalar field; it is symmetrical with respect to indices al , and evidently should be considered as the energy-momentum tensor for a scalar particle. For the current

$$S = 0 : \quad \nu^{(a)l}(x) = + \text{Sp} \left[\gamma^5 \gamma^l (+i A^* + \gamma^p A_p^*) \gamma^a (-i A + \gamma^s A_s) \right] , \quad (34)$$

an auxiliary table

$$\begin{array}{ccc} \gamma^5 \gamma^l & \gamma^a & \gamma^a \gamma^s \\ \gamma^5 \gamma^l \gamma^p & 0 & 0 \\ & 0 & \text{Sp} (\gamma^5 \gamma^l \gamma^p \gamma^a \gamma^s) \end{array} ,$$

and the current expression is

$$S = 0 : \quad \nu^{(a)l}(x) = i \epsilon^{alps} A_p^* A_s . \quad (35)$$

The current vanishes for all real-valued fields. And now consider the last current

$$S = 0 : \quad L^{(a)kl}(x) = + \text{Sp} \left[\sigma^{kl} (+i A^* + \gamma^p A_p^*) \gamma^a (-i A + \gamma^s A_s) \right] , \quad (36)$$

an auxiliary table

$$\begin{array}{ccc} \sigma^{kl} & \gamma^a & \gamma^a \gamma^s \\ \sigma^{kl} \gamma^p & 0 & \text{Sp} (\sigma^{kl} \gamma^a \gamma^s) \\ & \text{Sp} (\sigma^{kl} \gamma^p \gamma^a) & 0 \end{array} , \quad (37)$$

the non-zero traces

$$\frac{1}{4} \text{Sp} (\sigma^{kl} \gamma^a \gamma^s) = \frac{1}{2} (-g^{ka} g^{ls} + g^{ks} g^{la}) , \quad \frac{1}{4} \text{Sp} (\sigma^{kl} \gamma^s \gamma^a) = \frac{1}{2} (-g^{ks} g^{la} + g^{ka} g^{ls}) ,$$

an expression for the current is

$$S = 0 : \quad L^{(a)kl}(x) = \frac{i}{2} \left[(A^* A^k + A A^{k*}) g^{la} - (A^* A^l + A A^{l*}) g^{ka} \right] . \quad (38)$$

The current is non-zero both for complex- and real-valued field, it should be called the angular moment. Collecting together all results (take notice that we have no interpretation for the current $\nu^{(a)l}(x)$):

$S = 0$:

$$\begin{aligned} j^{(a)}(x) &= -i (A^* A^a - A A^{a*}) , \\ \nu^{(a)}(x) &\equiv 0 , \\ T^{(a)l}(x) &= A^* A g^{al} - g^{al} (A_s^* A^s) + (A^{a*} A^l + A^a A^{l*}) , \\ \nu^{(a)l}(x) &= i \epsilon^{alps} A_p^* A_s , \\ L^{(a)kl}(x) &= (i/2) [(A^* A^k + A A^{k*}) g^{la} - (A^* A^l + A A^{l*}) g^{ka}] . \end{aligned} \quad (39)$$

The same for pseudo scalar particle will look (and again, we have no interpretation for the current $\nu^{(a)l}(x)$)

$S = \tilde{0}$:

$$\begin{aligned} j^{(a)}(x) &= +i (B^* B^a - B B^{a*}) , \\ \nu^{(a)}(x) &\equiv 0 , \\ T^{(a)l}(x) &= B^* B g^{al} - g^{al} (B_s^* B^s) + (B^{a*} B^l + B^a B^{l*}) , \\ \nu^{(a)l}(x) &= -i \epsilon^{alps} B_p^* B_s , \\ L^{(a)kl}(x) &= (i/2) [(B^* B^k + B B^{k*}) g^{la} - (B^* B^l + B B^{l*}) g^{ka}] . \end{aligned} \quad (40)$$

Now let us consider the currents for a vector particle.

$$S = 1 : \quad j^{(a)}(x) = -\frac{1}{4} \text{Sp} \left[(+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+\gamma^s A_s + i \sigma^{mn} F_{mn}) \right] \quad (41)$$

an auxiliary table looks as

	$\gamma^a \gamma^s$	$\gamma^a \sigma^{mn}$
γ^p	0	$\text{Sp} (\gamma^p \gamma^a \sigma^{mn})$
σ^{bc}	$\text{Sp} (\sigma^{bc} \gamma^a \gamma^s)$	0

non-zero traces are

$$\begin{aligned} \frac{1}{4} \text{Sp} (\gamma^s \gamma^a \sigma^{mn}) &= \frac{1}{2} (-g^{ms} g^{na} + g^{ma} g^{ns}) , \\ \frac{1}{4} \text{Sp} (\gamma^a \gamma^s \sigma^{mn}) &= \frac{1}{2} (-g^{ma} g^{ns} + g^{ms} g^{na}) , \end{aligned}$$

so that

$$S = 1 : \quad j^{(a)} = +i (A_n^* F^{na} - A_n F^{na*}) \quad (42)$$

which vanishes for a real-valued field. For the current

$$S = 1 : \quad \nu^{(a)}(x) = -\frac{1}{4} \text{Sp} \left[\gamma^5 (+\gamma^p A_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+\gamma^s A_s + i \sigma^{mn} F_{mn}) \right] , \quad (43)$$

an auxiliary table

$$\begin{array}{ccc} \gamma^5 \gamma^p & \gamma^a \gamma^s & \gamma^a \sigma^{mn} \\ \gamma^5 \sigma^{bc} & 0 & \text{Sp} (\gamma^5 \gamma^p \gamma^a \sigma^{mn}) \\ & \text{Sp} (\gamma^5 \sigma^{bc} \gamma^a \gamma^s) & 0 \end{array} ,$$

non-zero traces

$$\frac{1}{4} \text{Sp} (\gamma^5 \gamma^p \gamma^a \sigma^{mn}) = \frac{i}{2} \epsilon^{pamn} , \quad \text{Sp} (\gamma^5 \sigma^{bc} \gamma^a \gamma^s) = \frac{i}{2} \epsilon^{asbc} ,$$

so that the current is

$$S = 1 : \quad \nu^{(a)} = -\frac{1}{2} (A_s^* F_{mn} - A_s F_{mn}^*) \epsilon^{asmn} . \quad (44)$$

It differs from zero only for complex-valued fields. Now for

$$S = 1 : \quad T^{(a)l}(x) = +\frac{1}{4} \text{Sp} [\gamma^l (+\gamma^p A_p^* + i\sigma^{bc} F_{bc}^*) \gamma^a (+\gamma^s A_s + i\sigma^{mn} F_{mn})] , \quad (45)$$

an auxiliary table looks

$$\begin{array}{ccc} \gamma^l \gamma^p & \gamma^a \gamma^s & \gamma^a \sigma^{mn} \\ \gamma^l \sigma^{bc} & \text{Sp} (\gamma^l \gamma^p \gamma^a \gamma^s) & 0 \\ & 0 & \text{Sp} (\gamma^l \sigma^{bc} \gamma^a \sigma^{mn}) \end{array} .$$

In order to obtain the formula for six-matrix product let us write down the formulas

$$\begin{aligned} \gamma^l \sigma^{bc} &= \frac{1}{2} (-\gamma^b g^{lc} + \gamma^c g^{lb} + i\gamma^5 \epsilon^{lbcd} \gamma_d) , \\ \gamma^a \sigma^{mn} &= \frac{1}{2} (-\gamma^m g^{an} + \gamma^n g^{am} + i\gamma^5 \epsilon^{amne} \gamma_e) , \end{aligned}$$

and then

$$\begin{aligned} (\gamma^l \sigma^{bc})(\gamma^a \sigma^{mn}) &= \frac{1}{4} (-\gamma^b g^{lc} + \gamma^c g^{lb} + i\gamma^5 \epsilon^{lbcd} \gamma_d) (-\gamma^m g^{an} + \gamma^n g^{am} + i\gamma^5 \epsilon^{amne} \gamma_e) = \\ &= \frac{1}{4} [-g^{lc} (-\gamma^b \gamma^m g^{an} + \gamma^b \gamma^n g^{am} + i\gamma^b \gamma^5 \epsilon^{amne} \gamma_e) + g^{lb} (-\gamma^c \gamma^m g^{an} + \gamma^c \gamma^n g^{am} + i\gamma^c \gamma^5 \epsilon^{amne} \gamma_e) + \\ &\quad + i\epsilon^{lbcd} (-\gamma^5 \gamma_d \gamma^m g^{an} + \gamma^5 \gamma_d \gamma^n g^{am} + i\gamma^5 \gamma_d \gamma^5 \epsilon^{amne} \gamma_e)] . \end{aligned}$$

Taking the trace of such a matrix expression we will arrive at the formula we need

$$\begin{aligned} &\frac{1}{4} \text{Sp} [(\gamma^l \sigma^{bc})(\gamma^a \sigma^{mn})] = \\ &= \frac{1}{4} [-g^{lc} (-g^{bm} g^{an} + g^{bn} g^{am}) + g^{lb} (-g^{cm} g^{an} + g^{cn} g^{am}) + \epsilon^{lbcd} \epsilon^{amne} g_{de}] . \end{aligned} \quad (46)$$

Now from (45) we get for the current

$$\begin{aligned} T^{(a)l} &= A_p^* A_s \frac{1}{4} \text{Sp} (\gamma^l \gamma^p \gamma^a \gamma^s) - F_{bc}^* F_{mn} \frac{1}{4} \text{Sp} [(\gamma^l \sigma^{bc})(\gamma^a \sigma^{mn})] = \\ &= A_p^* A_s (g^{lp} g^{as} - g^{la} g^{ps} + g^{ls} g^{pa}) - \\ &- \frac{1}{4} F_{bc}^* F_{mn} [-g^{lc}(-g^{bm} g^{an} + g^{bn} g^{am}) + g^{lb}(-g^{cm} g^{an} + g^{cn} g^{am}) + \epsilon^{lbcd} \epsilon^{amne} g_{de}] \end{aligned}$$

and further

$$T^{(a)l} = -g^{al} A_p^* A^p + (A^{a*} A^l + A^a A^{l*}) + F^{*ln} F_n^a - \frac{1}{4} F_{bc}^* F_{mn} \epsilon^{lbcd} \epsilon^{amne} g_{de} . \quad (47)$$

Here the last term may be changed to

$$- \frac{1}{4} F_{bc}^* F_{mn} \epsilon^{lbcd} \epsilon^{amne} g_{de} = + \frac{1}{4} F_{bc}^* F_{mn} \begin{vmatrix} g^{la} & g^{lm} & g^{ln} \\ g^{ba} & g^{bm} & g^{bn} \\ g^{ca} & g^{cm} & g^{cn} \end{vmatrix} = \frac{1}{2} F^{*mn} F_{mn} + F^{ln} F_n^{*a} .$$

With the help of that, from eq. (47) it follows

$$T^{(a)l} = -g^{al} A_p^* A^p + (A^{a*} A^l + A^a A^{l*}) + \frac{1}{2} F^{*mn} F_{mn} + (F^{*ln} F_n^a + F^{ln} F_n^{*a}) . \quad (48)$$

The tensor is symmetrical under indices al , it differs from zero for both complex- and real-valued fields, and may be considered as the energy-momentum tensor for a vector $S = 1$ particle. Now consider another current

$$S = 1 : \quad \nu^{(a)l}(x) = + \text{Sp} \left[\gamma^5 \gamma^l (+ \gamma^p A_p^* + i \sigma^{bc} F_{bc}^*) \gamma^a (+ \gamma^s A_s + i \sigma^{mn} F_{mn}) \right] , \quad (49)$$

an auxiliary table is

$$\begin{array}{ccc} \gamma^5 \gamma^l \gamma^p & \gamma^a \gamma^s & \gamma^a \sigma^{mn} \\ \gamma^5 \gamma^l \sigma^{bc} & \text{Sp} (\gamma^5 \gamma^l \gamma^p \gamma^a \gamma^s) & 0 \\ & 0 & \text{Sp} (\gamma^5 \gamma^l \sigma^{bc} \gamma^a \sigma^{mn}) \end{array} .$$

We will need the trace-formula for second term:

$$\begin{aligned} & \frac{1}{4} \text{Sp} [\gamma^5 (\gamma^l \sigma^{bc})(\gamma^a \sigma^{mn})] = \\ &= \frac{1}{16} \text{Sp} [\gamma^5 [-g^{lc}(-\gamma^b \gamma^m g^{an} + \gamma^b \gamma^n g^{am} + i \gamma^b \gamma^5 \epsilon^{amne} \gamma_e) + \\ & \quad + g^{lb}(-\gamma^c \gamma^m g^{an} + \gamma^c \gamma^n g^{am} + i \gamma^c \gamma^5 \epsilon^{amne} \gamma_e) + \\ & \quad + i \epsilon^{lbcd}(-\gamma^5 \gamma_d \gamma^m g^{an} + \gamma^5 \gamma_d \gamma^n g^{am} + i \gamma^5 \gamma_d \gamma^5 \epsilon^{amne} \gamma_e)] = \\ &= \frac{i}{4} [(g^{lc} \epsilon^{amnb} - g^{lb} \epsilon^{amnc}) + \epsilon^{lbcd} (\delta_d^n g^{am} - \delta_d^m g^{an})] . \end{aligned}$$

with the help of that, for the current we will find

$$\nu^{(a)l}(x) = i A_p^* A_s \epsilon^{alps} + \frac{i}{2} F_{bc}^* F_{mn} [g^{an} \epsilon^{lbcm} + g^{lb} \epsilon^{amnc}] . \quad (50)$$

The conservation laws for the quantity $\nu^{(a)l}(x)$ may be written down in detailed form:

$$\begin{aligned} \nu^{(a)0} : \quad & \nu^{(0)0} = -i [+E^{*1} B^1 + E^{*2} B^2 + E^{*3} B^3 - \text{c.c}] , \\ & \nu^{(1)0} = +i [-A_2^* A_3 + (E_2^* E_3 + B_2^* B_3) - \text{c.c}] , \\ & \nu^{(2)0} = +i [-A_3^* A_1 + (E_3^* E_1 + B_3^* B_1) - \text{c.c}] , \\ & \nu^{(3)0} = +i [-A_1^* A_2 + (E_1^* E_2 + B_1^* B_2) - \text{c.c}] , \end{aligned} \quad (51)$$

$$\begin{aligned} \nu^{(a)1} : \quad & \nu^{(0)1}(x) = +i [+A_2^* A_3 + (E_2^* E_3 + B_2^* B_3) - \text{c.c}] , \\ & \nu^{(1)1}(x) = -i [-E^{*1} B^1 + E^{*2} B^2 + E^{*3} B^3 - \text{c.c}] , \\ & \nu^{(2)1}(x) = i [-A_0^* A_3 + (E_2^* B_1 + E_1^* B_2) - \text{c.c}] , \\ & \nu^{(3)1}(x) = i [+A_0^* A_2 + (E_3^* B_1 + E_1^* B_3) - \text{c.c}] , \end{aligned} \quad (52)$$

$$\begin{aligned} \nu^{(a)2} : \quad & \nu^{(0)2}(x) = +i [+A_3^* A_1 + (E_3^* E_1 + B_3^* B_1) - \text{c.c}] , \\ & \nu^{(1)2}(x) = i [+A_0^* A_3 + (E_1^* B_2 + E_2^* B_1) - \text{c.c}] , \\ & \nu^{(2)2}(x) = -i [+E^{*1} B^1 - E^{*2} B^2 + E^{*3} B^3 - \text{c.c}] , \\ & \nu^{(3)2}(x) = i [-A_0^* A_1 + (E_3^* B_2 + E_2^* B_3) - \text{c.c}] , \end{aligned} \quad (53)$$

$$\begin{aligned} \nu^{(a)3} : \quad & \nu^{(0)3}(x) = +i [+A_1^* A_2 + (E_1^* E_2 + B_1^* B_2) - \text{c.c}] , \\ & \nu^{(1)3}(x) = i [-A_0^* A_2 + (E_1^* B_3 + E_3^* B_1) - \text{c.c}] , \\ & \nu^{(2)3}(x) = i [+A_0^* A_1 + (E_2^* B_3 + E_3^* B_2) - \text{c.c}] , \\ & \nu^{(3)3}(x) = -i [+E^{*1} B^1 + E^{*2} B^2 - E^{*3} B^3 - \text{c.c}] . \end{aligned} \quad (54)$$

And finally, let us obtain an explicit form of the current

$$S = 1 : \quad L^{(a)kl}(x) = +\frac{1}{4} \text{Sp}; \left[\sigma^{kl} (+\gamma^p A_p^* + i\sigma^{bc} F_{bc}^*) \gamma^a (+\gamma^s A_s + i\sigma^{mn} F_{mn}) \right] , \quad (55)$$

an auxiliary table is

$$\begin{array}{ccc} \sigma^{kl} \gamma^p & \gamma^a \gamma^s & \gamma^a \sigma^{mn} \\ \sigma^{kl} \sigma^{bc} & 0 & \text{Sp} (\sigma^{kl} \gamma^p \gamma^a \sigma^{mn}) \\ & \text{Sp} (\sigma^{kl} \sigma^{bc} \gamma^a \gamma^s) & 0 \end{array} ,$$

non-zero traces are

$$\begin{aligned} & \frac{1}{4} \text{Sp} (\sigma^{kl} \sigma^{bc} \gamma^a \gamma^s) = \frac{1}{4} \text{Sp} (\sigma^{bc} \gamma^a \gamma^s \sigma^{kl}) = \\ & = \frac{1}{16} \text{Sp} [(\gamma^b g^{ca} - \gamma^c g^{ba} + i\gamma^5 \epsilon^{bcad} \gamma_d)(-\gamma^k g^{sl} + \gamma^l g^{sk} + i\gamma^5 \epsilon^{skle} \gamma_e)] = \\ & = \frac{1}{4} [g^{ca}(-g^{bk} g^{sl} + g^{bl} g^{sk}) - g^{ba}(-g^{ck} g^{sl} + g^{cl} g^{sk}) - \epsilon^{bcad} \epsilon^{skle} g_{de}] \\ & \frac{1}{4} \text{Sp} (\sigma^{kl} \gamma^p \gamma^a \sigma^{mn}) = \\ & = \frac{1}{16} \text{Sp} [(\gamma^k g^{lp} - \gamma^l g^{kp} + i\gamma^5 \epsilon^{klpd} \gamma_d)(-\gamma^m g^{an} + \gamma^n g^{am} + i\gamma^5 \epsilon^{amne} \gamma_e)] = \\ & = \frac{1}{4} [g^{lp}(-g^{km} g^{an} + g^{kn} g^{am}) - g^{kp}(-g^{lm} g^{an} + g^{ln} g^{am}) + \epsilon^{klpd} \epsilon^{amne} g_{de}] . \end{aligned}$$

With the help of two relations

$$\begin{aligned}
& i F_{bc}^* A_s \frac{1}{4} \text{Sp} (\sigma^{kl} \sigma^{bc} \gamma^a \gamma^s) = \\
& = \frac{i}{2} \left[-A^a F^{*kl} + (F^{*ak} A^l - F^{*al} A^k) + (-g^{ak} F^{*ls} A_s + g^{al} F^{*ks} A_s) \right] , \\
& i A_p^* F_{mn} \frac{1}{4} \text{Sp} (\sigma^{kl} \gamma^p \gamma^a \sigma^{mn}) = \\
& \frac{i}{2} \left[-A^{*a} F^{kl} + (F^{ak} A^{*l} - F^{al} A^{*k}) + (-g^{ak} F^{ls} A_s^* + g^{al} F^{ks} A_s^*) \right]
\end{aligned}$$

one readily produces the expression for this current

$$\begin{aligned}
L^{(a)kl} = \frac{i}{2} \{ & -(A^a F^{*kl} + A^{*a} F^{kl}) + [(F^{*ak} A^l - F^{*al} A^k) + (F^{ak} A^{*l} - F^{al} A^{*k})] + \\
& + [(-g^{ak} F^{*ls} A_s + g^{al} F^{*ks} A_s) + (-g^{ak} F^{ls} A_s^* + g^{al} F^{ks} A_s^*)] \} . \quad (56)
\end{aligned}$$

Collecting together all results for a particle of $S = 1$:

$$\begin{aligned}
S = 1 : \quad & j^{(a)} = +i (A_n^* F^{na} - A_n F^{na*}) , \\
& \nu^{(a)} = -(1/2) (A_s^* F_{mn} - A_s F_{mn}^*) \epsilon^{asmn} , \\
& T^{(a)l} = -g^{al} A_p^* A^p + (A^{a*} A^l + A^a A^{l*}) + \\
& + (1/2) F^{*mn} F_{mn} + (F^{*ln} F_n^a + F^{ln} F_n^{*a}) , \\
& \nu^{(a)l}(x) = i A_p^* A_s \epsilon^{alps} + \\
& + (i/2) F_{bc}^* F_{mn} (g^{an} \epsilon^{lbcm} + g^{lb} \epsilon^{amnc}) , \\
& L^{(a)kl} = (i/2) \{ -(A^a F^{*kl} + A^{*a} F^{kl}) + \\
& + [(F^{*ak} A^l - F^{*al} A^k) + (F^{ak} A^{*l} - F^{al} A^{*k})] + \\
& + [(-g^{ak} F^{*ls} A_s + g^{al} F^{*ks} A_s) + (-g^{ak} F^{ls} A_s^* + g^{al} F^{ks} A_s^*)] \} . \quad (57)
\end{aligned}$$

Take notice that we have no interpretation for two currents, $\nu^{(a)}$ and $\nu^{(a)l}(x)$, which are non-zero for complex-valued fields only. Here we will not consider in detail the case of vector particle with another intrinsic parity, particle of the type $S = \tilde{1}$, any additional ideas will not arise at this.

References

- [1] Darvin C.G. Proc. Roy. Soc. Lond. A. **118**, No 780. 654-680 (1928).
- [2] Ivanenko D, Landau L. Z. Phys. **48**. 340-348 (1928).
- [3] Kähler E. Abh. Dt. Akad. Wiss. Berlin, Kl. Math.-Phys. u. Techn. No 4 (1960); No 1 (1961).

- [4] Kähler E. Rendiconti di Matematica. (Roma). **21**. 425-523 (1962).
- [5] Durand E. Phys. Rev. **D11**. 3405-3416 (1975).
- [6] Strajev V.I. Izvestia Vuzov, Fizika. No 8. 45-48 (1977).
- [7] Pestov A. B. TMF. **34**. 48-57 (1978).
- [8] Graf Wolfgang. Annales Inst. H. Poincaré. A. **29**. 85-109 (1978).
- [9] Strajev V.I. Acta Phys. Polon. **B9**. 449-458 (1978).
- [10] Bogush A.A., Kruglov C.I., Strajev V.I. Doklady AN BSSR. **22**. 893-895 (1978).
- [11] Bogush A.A., Kruglov C.I. Vesti AN BSSR. ser. fiz.-mat. **4**. 58-65. (1978).
- [12] Benn I.M., Tucker R.W. Phys. Lett. **B119**. 348-350 (1982); Commun. Math. Phys. **89**. 341-362 (1983); J. Phys. **A16**. 4147-4153 (1983); Phys. Lett. **B130**. 177-178 (1983); Phys. Lett. **B132**. 325-327 (1983).
- [13] Ivanenko D.D., Obukhov Yu.N., Solodukhin S.N. On antisymmetric tensor representation of the Dirac equation. – Miramare-Trieste, 1985. (Preprint / Miramare-Trieste ; IC/85/2).
- [14] Bullinaria J.A. Ann. Phys. (N.Y.) **168**. 301-343 (1986).
- [15] Drechsler W., Thacker W. Class. Quantum Grav. **4**. 291-318 (1987).
- [16] Satikov I.A., Strajev V.I. TMF. **73**. 16-25 (1987).
- [17] Krolikowski W. Acta Phys. Polon. **B20**. 849-858 (1989); **B21**. 201-207 (1990).
- [18] Pletjuxov V.A., Strajev V.I. Doclady AN BSSR. **33**. 328-331 (1989);
- [19] Pletjuxov V.A., Strajev V.I. Yadernaya Fizika. **49**. 1505-1514 (1989).
- [20] Redkov V.M. Vesti AN BSSR. ser. fiz.-mat. No 1. 90-95 (2000).